ANATOLIA FEED

nektar

EC FERTILIZER
NK GÜBRESİ HARMANLANMIŞ

ANATOLIA FEED

10-0-46

GARANTİ EDİLEN İÇERİĞİ	%(w/w)
Toplam Azot (N)	10
Üre Azotu (NH_2-N)	10
Suda Çözünür Potasyum Oksit (K_2O)	46

NET :

ANATOLIA FEED

ANATOLIA FEED

EC FERTILIZER
NK GÜBRESİ HARMANLANMIŞ

ANATOLIA FEED

10-0-46

UYGULAMA DOZAJLARI KULLANIM DÖNEMLERİ

15-30-15+ME / 15-30-15+TE

ÜRÜN	UYGULAMA DÖNEMİ	UYGULAMA DOZU YAPRAKTAN	UYGULAMA DOZU TOPRAKTAN
NARENCİYE	Çiçeklenme öncesi	200-250 g/100 L suyu	2-4 kg/da
ELMA, ARMUT VB.	Çiçeklenme öncesi	200-250 g/100 L suyu	2-4 kg/da
ŞEFTALİ, KAYISI, KİRAZ, VİŞNE, ERİK ve NAR	Çiçeklenme öncesi	200-250 g/100 L suyu	2-4 kg/da
BAĞ	Filizlenme başlangıcı	200-250 g/100 L suyu	1-3 kg/da
ANTEP FISTIĞI, FINDIK	Çiçeklenme öncesi	200-250 g/100 L suyu	2-4 kg/da
ZEYTİN	Çiçeklenme öncesi	200-250 g/100 L suyu	2-4 kg/da
MUZ	Doğum öncesi 15 gün ara ile 2 uygulama	200 g/100 L suyu	1-3 kg/da
KİVİ, İNCİR	Çiçeklenme öncesi	200-250 g/100 L suyu	2-4 kg/da
DOMATES, BİBER	Çiçeklenme dönemi 15 gün ara ile 3 uygulama	200-250 g/100 L suyu	1-3 kg/da
HIYAR, PATLICAN	Çiçeklenme dönemi 15 gün ara ile 3 uygulama	200-250 g/100 L suyu	1-3 kg/da
KAVUN, KAPUZ	Çiçeklenme dönemi 15 gün ara ile 3 uygulama	200-250 g/100 L suyu	1-3 kg/da
KESME ÇİÇEKÇİLİK	Gelişim dönemi boyunca 3 uygulama	200-250 g/100 L suyu	1-3 kg/da
MERCİMEK, NOHUT	Çiçeklenme dönemi	200-250 g/da	-
YERFİSTİĞİ	Çiçeklenme dönemi	200-250 g/100 L suyu	1-3 kg/da
PAMUK, AYÇİÇEĞİ, MISIR, KANOLA, TÜTÜN vb.	Bitkiler 20-30 cm iken 15 gün ara ile 2 uygulama	200-250 g/da	2-4 kg/da
ÇİLEK	Karakşlenme dönemi	200-300 g/da	-
ÇAY	Gelişim dönemi boyunca 2 uygulama	200-250 g/100 L suyu	1-2 kg/da
ŞEKER PANCARI	Yumru oluşumu dönemi 2 uygulama	200-250 g/100 L suyu	2-3 kg/da
MEYVE FİDANLARI	Gelişim dönemi boyunca 3 uygulama	250-300 g/100 L suyu	2-4 kg/da
HUBUBAT	Karakşlenme dönemi	200-300 g/da	-
YEŞİL ALAN BİTKİLERİ	Gelişim dönemi boyunca	200-250 g/da	1-3 kg/da

15-5-30+TE / 15-5-30+ME / 16-8-24+ME / 16-8-24+TE / 3-5-40+ME / 3-5-40+TE / 10-0-46

ÜRÜN	UYGULAMA DÖNEMİ	UYGULAMA DOZU YAPRAKTAN	UYGULAMA DOZU TOPRAKTAN
NARENCİYE	Meyve büyüme dönemi 2 uygulama	250-300 g/100 L suyu	2-4 kg/da
ELMA, ARMUT VB.	Meyve büyüme dönemi 2 uygulama	250-300 g/100 L suyu	2-4 kg/da
ŞEFTALİ, KAYISI, KİRAZ, VİŞNE, ERİK ve NAR	Meyve büyüme dönemi 2 uygulama	250-300 g/100 L suyu	2-4 kg/da
BAĞ	Tiafil su çıkıma dönemi 2 uygulama	200-250 g/100 L suyu	1-3kg/da
ANTEP FISTIĞI, FINDIK	Tane dolum dönemi	250-300 g/100 L suyu	2-4 kg/da
ZEYTİN	Tane irlme dönemi	200-250 g/100 L suyu	2-4 kg/da
MUZ	Meyve şişirme dönemi	200-250 g/100 L suyu	1-3 kg/da
KİVİ, İNCİR	Meyve büyüme dönemi	200-250 g/100 L suyu	2-4 kg/da
DOMATES, BİBER	Meyve büyüme dönemi	200-250 g/100 L suyu	1-3 kg/da
HIYAR, PATLICAN	Meyve büyüme dönemi	200-250 g/100 L suyu	1-3 kg/da
KAVUN, KAPUZ	Meyve büyüme dönemi	200-250 g/100 L suyu	1-3 kg/da
KESME ÇİÇEKÇİLİK	Gelişim dönemi boyunca	250-300 g/100 L suyu	1-3 kg/da
MERCİMEK, NOHUT	Tane dolum dönemi	200-250 g/100 L suyu	-
YERFİSTİĞİ	İç dolum dönemi	250-300 g/100 L suyu	1-3 kg/da
PAMUK, AYÇİÇEĞİ, MISIR, KANOLA, TÜTÜN vb.	Bitkiler 40-50 cm iken 15 gün ara ile 2 uygulama	250-300 g/100 L suyu	2-4 kg/da
ÇİLİK	Tane dolum dönemi	200-250 g/100 L suyu	-
ÇAY	Gelişim dönemi boyunca	200-250 g/100 L suyu	1-3 kg/da
ŞEKER PANCARI	Yumru büyüme dönemi	200-250 g/100 L suyu	2-4 kg/da
MEYVE FİDANLARI	Gelişim dönemi boyunca	200-250 g/100 L suyu	1-3 kg/da
HUBUBAT	Tane dolum dönemi	200-250 g/100 L suyu	-
YEŞİL ALAN BİTKİLERİ	Gelişim dönemi	200-250 g/100 L suyu	1-3 kg/da

18-18-18+ME / 18-18-18+TE / 20-10-20+ME / 20-10-20+TE / 20-20-20 / 30-5 -0

ÜRÜN	UYGULAMA DÖNEMİ	UYGULAMA DOZU YAPRAKTAN	UYGULAMA DOZU TOPRAKTAN
NARENCİYE	Filizlenme dönemi	250-300 g/100 L suyu	2-4 kg/da
ELMA, ARMUT VB.	Gözler uyanma dönemi	250-300 g/100 L suyu	2-4 kg/da
ŞEFTALİ, KAYISI, KİRAZ, VİŞNE, ERİK ve NAR	Gözler uyanma dönemi	250-300 g/100 L suyu	2-4 kg/da
BAĞ	Gözler uyanma dönemi	250-300 g/100 L suyu	2-4 kg/da
ANTEP FISTIĞI, FINDIK	Gözler uyanma dönemi	200-250 g/100 L suyu	1-3 kg/da
ZEYTİN	Gözler uyanma dönemi	250-300 g/100 L suyu	2-4 kg/da
MUZ	Fide dikiminden itibaren 15 gün ara ile 3 uygulama	250-300 g/100 L suyu	2-4 kg/da
KİVİ, İNCİR	Gözler uyanma evresi	250-300 g/100 L suyu	2-4 kg/da
DOMATES, BİBER	Gelişim dönemi boyunca 3 uygulama	200-250 g/100 L suyu	1-3 kg/da
HIYAR, PATLICAN	Gelişim dönemi boyunca 3 uygulama	200-250 g/100 L suyu	1-3 kg/da
KAVUN, KAPUZ	Gelişim dönemi boyunca 3 uygulama	200-250 g/100 L suyu	1-3 kg/da
KESME ÇİÇEKÇİLİK	Gelişim dönemi boyunca 2 uygulama	200-250 g/100 L suyu	1-3 kg/da
MERCİMEK, NOHUT	Çimlenme dönemnden sonra 2 uygulama	200-250 g/100 L suyu	-
YERFİSTİĞİ	Çimlenme dönemnden sonra 2 uygulama	200-250 g/100 L suyu	1-3 kg/da
PAMUK, AYÇİÇEĞİ, MISIR, KANOLA, TÜTÜN vb.	Bitkiler 10-20 cm iken	200-250 g/100 L suyu	-
ÇİLİK	Sapa kalkma dönemi	200-250 g/100 L suyu	1-3 kg/da
ÇAY	Gelişim dönemi boyunca	200-250 g/100 L suyu	1-3 kg/da
ŞEKER PANCARI	Yumru gelişim dönemi 2 uygulama	200-300 g/100 L suyu	2-4 kg/da
MEYVE FİDANLARI	Gelişim dönemi boyunca 2 uygulama	200-300 g/100 L suyu	1-3 kg/da
HUBUBAT	Sapa kalkma dönemi	200-250 g/100 L suyu	3-5 kg/da
YEŞİL ALAN BİTKİLERİ	Gelişim dönemi boyunca 2 uygulama	200-250 g/100 L suyu	1-3 kg/da

APPLICATION DOSES AND USAGE PERIODS

15-30-15+ME / 15-30-15+TE

CROP	APPLICATION PERIOD	APPLICATION DOSAGE FOLIAR	APPLICATION DOSAGE SOIL
CITRUS	Before Flowering	200-250 g/100 L water	2-4 kg/da
APPLE, PEAR ETC	Before Flowering	200-250 g/100 L water	2-4 kg/da
PEACH, APRICOT, CHERRY, SOUR CHERRY, PLUM and POMEGRANATE	Before Flowering	200-250 g/100 L water	2-4 kg/da
VINEYARD	Beginning of shooting	200-250 g/100 L water	1-3 kg/da
PISTACHIO, HAZELNUT	Before Flowering	200-250 g/100 L water	2-4 kg/da
OLIVE	Before Flowering	200-250 g/100 L water	2-4 kg/da
BANANA	2 applications with 15 days interval before bunch emergence	200 g/100 L water	1-3 kg/da
KIWI, FIG	Before Flowering	200-250 g/100 L water	2-4 kg/da
TOMATO, PEPPER	Flowering period 3 applications with 15 days intervals	200-250 g/100 L water	1-3 kg/da
CUCUMBER, EGGPLANT	Flowering period 3 applications with 15 days intervals	200-250 g/100 L water	1-3 kg/da
MELON, WATERMELON	Flowering period 3 applications with 15 days intervals	200-250 g/100 L water	1-3 kg/da
CUT FLOWER	3 applications during growth period	200-250 g/100 L water	1-3 kg/da
LENTIL, CHICKPEA	Flowering period	200-250 g/da	-
PEANUT	Flowering period	200-250 g/100 L water	1-3 kg/da
COTTON, SUNFLOWER, CORN, CANOLA, TOBACCO ETC.	2 applications with 15 days interval when the plants are 20-30 cm	200-250 g/da	2-4 kg/da
RICE	Tillering period	200-300 g/da	-
TEA PLANT	2 applications during growth period	200-250 g/100 L water	1-2 kg/da
SUGAR BEET	2 applications tuber formation	200-250 g/100 L water	2-3 kg/da
FRUIT SAPLING	2 applications during growth period	250-300 g/100 L water	2-4 kg/da
CEREALS	Tillering period	200-300 g/da	-
GREEN AREAS	During growth period	200-250 g/da	1-3 kg/da

15-5-30+TE / 15-5-30+ME / 16-8-24+ME / 16-8-24+TE / 3-5-40+ME / 3-5-40+TE / 10-0-46

CROP	APPLICATION PERIOD	APPLICATION DOSAGE FOLIAR	APPLICATION DOSAGE SOIL
CITRUS	Fruit growth period 2 applications	250-300 g/ 100 L water	2-4 kg/da
APPLE, PEAR ETC	Fruit growth period 2 applications	250-300 g/ 100 L water	2-4 kg/da
PEACH, APRICOT, CHERRY, SOUR CHERRY, PLUM and POMEGRANATE	Fruit growth period 2 applications	250-300 g/ 100 L water	2-4 kg/da
VINEYARD	Veraison period 2 applications	200-250 g/100 L water	1-3 kg/da
PISTACHIO, HAZELNUT	Grain Growth Period	250-300 g/100 L water	2-4 kg/da
OLIVE	Grain Ripening Period	200-250 g/100 L water	2-4 kg/da
BANANA	Fruit Growth Period	200-250 g/100 L water	1-3 kg/da
KIWI, FIG	Fruit Growth Period	200-250 g/100 L water	2-4 kg/da
TOMATO, PEPPER	Fruit Growth Period	200-250 g/100 L water	1-3 kg/da
CUCUMBER, EGGPLANT	Fruit Growth Period	200-250 g/100 L water	1-3 kg/da
MELON, WATERMELON	Fruit Growth Period	200-250 g/100 L water	1-3 kg/da
CUT FLOWER	During Growth Period	250-300 g/100 L water	1-3 kg/da
LENTIL, CHICKPEA	Grain Growth Period	200-250 g/100 L water	-
PEANUT	Internal Filling Period	250-300 g/100 L water	1-3 kg/da
COTTON, SUNFLOWER, CORN, CANOLA, TOBACCO ETC.	2 applications with 15 days interval when the plants are 40-50 cm	250-300 g/100 L water	2-4 kg/da
RICE	Grain Growth Period	200-250 g/100 L water	-
TEA PLANT	During Growth Period	200-250 g/100 L water	1-3 kg/da
SUGAR BEET	Tuber Growth Period	200-250 g/100 L water	2-4 kg/da
FRUIT SAPLING	During Growth Period	200-250 g/100 L water	1-3 kg/da
CEREALS	Grain Growth Period	200-250 g/100 L water	-
GREEN AREAS	Growth Period	200-250 g/100 L water	1-3 kg/da

18-18-18+ME / 18-18-18+TE / 20-10-20+ME / 20-10-20+TE / 20-20-20 / 30-5 -0

CROP	APPLICATION PERIOD	APPLICATION DOSAGE FOLIAR	APPLICATION DOSAGE SOIL
CITRUS	Shooting Period	250-300 g/100 L water	2-4 kg/da
APPLE, PEAR ETC	Budbreak Period	250-300 g/100 L water	2-4 kg/da
PEACH, APRICOT, CHERRY, SOUR CHERRY, PLUM and POMEGRANATE	Budbreak Period	250-300 g/100 L water	2-4 kg/da
VINEYARD	Budbreak Period	250-300 g/100 L water	2-4 kg/da
PISTACHIO, HAZELNUT	Budbreak Period	200-250 g/100 L water	1-3 kg/da
OLIVE	Budbreak Period	250-300 g/100 L water	2-4 kg/da
BANANA	3 applications with an interval of 15 days from Budbreak Period	250-300 g/100 L water	2-4 kg/da
KIWI, FIG	Budbreak Period	250-300 g/100 L water	2-4 kg/da
TOMATO, PEPPER	3 applications during the growth period	200-250 g/100 L water	1-3 kg/da
CUCUMBER, EGGPLANT	3 applications during the growth period	200-250 g/100 L water	1-3 kg/da
MELON, WATERMELON	3 applications during the growth period	200-250 g/100 L water	1-3 kg/da
CUT FLOWER	2 applications during the growth period	200-250 g/100 L water	1-3 kg/da
LENTIL, CHICKPEA	2 applications after the germination period	200-250 g/100 L water	-
PEANUT	2 applications after the germination period	200-250 g/100 L water	1-3 kg/da
COTTON, SUNFLOWER, CORN, CANOLA, TOBACCO ETC.	When plants 10-20 cm	200-250 g/100 L water	-
RICE	Stem elongation period	200-250 g/100 L water	1-3 kg/da
TEA PLANT	During growth period	200-250 g/100 L water	1-3 kg/da
SUGAR BEET	2 applications tuber growth period	250-300 g/100 L water	2-4 kg/da
FRUIT SAPLING	2 applications during the growth period	200-300 g/100 L water	1-3 kg/da
CEREALS	Stem elongation period	200-250 g/100 L water	3-5 kg/da
GREEN AREAS	2 applications during the growth period	200-250 g/100 L water	1-3 kg/da

UYARILAR : * Dikkat çocuklardan, evcil hayvanlardan ve gıdalardan uzak tutunuz. * Asla uygun doz oranlarını aşmayınız ve sadece gerekli olan yerlerde kullanınız. * Göz, yüz ve deri ile temas ettirmeyiniz. *Temas halinde bol su ile yıkayınız. * Önerilen dozajı aşmayınız.* Kullanmdan sonra ellerinizi bol su ile yıkayınız. * Kullanılmış olan boş ambalajları başka amaçlarla kullanmayınız.* Kullandıktan sonra ambalajları imha ediniz. *Toprak ve / veya yaprak analizi yaptırılarak gübre kullanılması tavsiye edilir.

DEPOLAMA :* Ürün, kendi orijinal ambalajında, serin ve kuru ortamda **0 °C / + 45 °C** arasında, depolandığında, fiziksel ve kimyasal özelliklerini en az **3 yıl** korur.* Bu nederle, Ürünü kendi orijinal ambalajında, kuru ve serin bir yerde, belirtilen sıcaklıklar arasında muhafaza ediniz. * Ürünü, her ne şekilde olursa olsun, güneş ışığına maruz bırakmayınız.

FİRMA BEYANI : İmalatçı firma, tavsiyelerine uyularak kullanıldığında, ürünün kalitesini garanti eder. Ancak, üretici tarafından, ürünün kullanımı, firmamızın kontrolü dışında olduğundan, hatalı kullanım ve karışımlardan doğuş okuşabilecek olumsuzluklar ile amaç dişı kullanım ve depolamadan doğacak sorunlardan firmamız mesuliyet kabul etmez. Ürünün, spesifikasyonuna uygunluğu, sadece,ürünün teslimi sırasında, firma garantisi altındadır.

Piyasaya sunacağımız etiket ile aynı olacağını taahhüt eder, aksinin ispatı halinde, 7223 saylı kanuna göre, hakkımızda yapılacak cezai işlemleri, peşinen kabul ederiz.

ASLA UYGUN DOZ ORANLARINI AŞMAYINIZ!
SADECE GEREKLİ YERLERDE KULLANINIZ!

Tescil No / Reg. No :
Lisans No / Lic. No : 116
Üretim Tarihi / Prod. Date :
Son Kullanma Tarihi / Exp. Date:
Parti No / Lot No :

WARNING: • Keep away from children, animals and food • Do not exceed the recommended dose and use only where necessary• Avoid contact with eyes, face and skin. • **Was with plenty of water in contact** • Wash hands with plenty of water after use. • **Do not exceed recommended dosage** • Do not use empty packages for any other purposes • Destroy empty packages after use. • It is recommended to make an analysis of the soil before application of the fertilizer.

STORAGE: When stored in the original packaging, in cool and dry area, in the temperature between **0–45 °C**, the product keeps its physical and chemical characteristics for **3 years.** **For this reason, keep the product its original packages; in dry and cool area and between indicated temperatures.** Keep it away from direct sunlight.

COMPANY STATEMENT: The manufacturer company guarantees the quality of the product when used according to recommended methods of application. **The comformity of the product to its specification is only guaranteed by the company at the time of delivery of product.** The product is out of warranty and the manufacturer is not responsible for all damage if it is stored out of cautions or used out of recommendation

ÜRETİCİ FİRMA

nektar

NEKTAR TARIM TIC. PAZ. LTD. ŞTİ.
AOSB 2. Kısım Mahallesi 24. Cadde No:2 Döşemealtı / ANTALYA
Tel: 0 242 258 01 41-47 Faks: 0 242 258 01 48
info@nektartarim.com / www.nektartarim.com

ANATOLIA FEED

ANATOLIA FEED